

Original Paper

Development and Feasibility Testing of Internet-Delivered Acceptance and Commitment Therapy for Severe Health Anxiety: Pilot Study

Ditte Hoffmann¹, MSc; Charlotte Ulrikka Rask^{1,2}, MD, PhD; Erik Hedman-Lagerlöf³, MSc, PhD; Brjánn Ljótsson^{3,4}, MSc, PhD; Lisbeth Frostholt¹, MSc, PhD

¹The Research Clinic for Functional Disorders and Psychosomatics, Aarhus University Hospital, Aarhus, Denmark

²Child and Adolescent Psychiatric Centre, Aarhus University Hospital, Aarhus, Denmark

³Department of Clinical Neuroscience, Karolinska Institute, Stockholm, Sweden

⁴Centre for Psychiatry Research, Karolinska Institute, Stockholm, Sweden

Corresponding Author:

Ditte Hoffmann, MSc

The Research Clinic for Functional Disorders and Psychosomatics

Aarhus University Hospital

Noerrebrogade 44

Bldg 4, 1

Aarhus, 8000

Denmark

Phone: 45 78464625

Fax: 45 78464340

Email: dittjese@rm.dk

Abstract

Background: Severe health anxiety (hypochondriasis), or illness anxiety disorder according to the Diagnostic and Statistical Manual of Mental Disorders, 5th edition, is characterized by preoccupation with fear of suffering from a serious illness in spite of medical reassurance. It is a debilitating, prevalent disorder associated with increased health care utilization. Still, there is a lack of easily accessible specialized treatment for severe health anxiety.

Objective: The aims of this paper were to (1) describe the development and setup of a new internet-delivered acceptance and commitment therapy (iACT) program for patients with severe health anxiety using self-referral and a video-based assessment; and (2) examine the feasibility and potential clinical efficacy of iACT for severe health anxiety.

Methods: Self-referred patients (N=15) with severe health anxiety were diagnostically assessed by a video-based interview. They received 7 sessions of clinician-supported iACT comprising self-help texts, video clips, audio files, and worksheets over 12 weeks. Self-report questionnaires were obtained at baseline, post-treatment, and at 3-month follow-up. The primary outcome was Whiteley-7 Index (WI-7) measuring health anxiety severity. Depressive symptoms, health-related quality of life (HRQoL), life satisfaction, and psychological flexibility were also assessed. A within-group design was employed. Means, standard deviations, and effect sizes using the standardized response mean (SRM) were estimated. Post-treatment interviews were conducted to evaluate the patient experience of the usability and acceptability of the treatment setup and program.

Results: The self-referral and video-based assessments were well received. Most patients (12/15, 80%) completed the treatment, and only 1 (1/15, 7%) dropped out. Post-treatment (14/15, 93%) and 3-month follow-up (12/15, 80%) data were available for almost all patients. Paired t tests showed significant improvements on all outcome measures both at post-treatment and 3-month follow-up, except on one physical component subscale of HRQoL. Health anxiety symptoms decreased with 33.9 points at 3-month follow-up (95% CI 13.6-54.3, $t_{11} = 3.66$, $P = .004$) with a large within-group effect size of 1.06 as measured by the SRM.

Conclusions: Treatment adherence and potential efficacy suggest that iACT may be a feasible treatment for health anxiety. The uncontrolled design and small sample size of the study limited the robustness of the findings. Therefore, the findings should be replicated in a randomized controlled trial. Potentially, iACT may increase availability and accessibility of specialized treatment for health anxiety.

Trial Registration: Danish Data Protection Agency, Central Denmark Region: 1-16-02-427-14; <https://www.rm.dk/sundhed/faginfo/forskning/datatilsynet/> (Archived by Webcite at <http://www.webcitation.org/6yDA7WovM>)

(*JMIR Ment Health* 2018;5(2):e28) doi:[10.2196/mental.9198](https://doi.org/10.2196/mental.9198)

KEYWORDS

health anxiety; illness anxiety disorder; hypochondriasis; internet intervention; feasibility; acceptance and commitment therapy; behavior therapy

Introduction

Severe health anxiety is characterized by the preoccupation with fear of having a serious illness, which interferes with daily functions and persists despite medical reassurance [1]. In the Diagnostic and Statistical Manual of Mental Disorders, 4th edition (DSM-IV), the disorder was labeled hypochondriasis, but in the new Diagnostic and Statistical Manual of Mental Disorders, 5th edition (DSM-5) [2], severe health anxiety is now classified as illness anxiety disorder or somatic symptom disorder, if persistent and distressing physical symptoms are present. Severe health anxiety is highly prevalent in medical settings (0.3% to 8.5%) [1,3-5], and has an estimated lifetime prevalence of 5.7% in the general population [6].

Despite the high prevalence and the existence of effective psychological treatments [7], the disorder is still rarely diagnosed in the health care setting [8]. Health care professionals often lack the necessary knowledge about health anxiety, and when the disorder is recognized, there is often a lack of proper treatment options. Furthermore, a subgroup of patients tends to be care-avoidant, and this group is therefore at an even higher risk of diagnostic delay. As severe health anxiety is associated with extensive use of health care services [9,10] and occupational disability [11], and rarely remits if untreated [10], the limited accessibility to proper assessment and treatment is a major challenge to the health care system. Thus, evidence-based, easily accessible treatment options are warranted for this debilitating disorder.

Patient self-referral in combination with internet-delivered psychological treatment is a new approach that may facilitate treatment accessibility and broaden availability. As used here, internet-delivered treatment is a highly structured treatment that the patient accesses through a secure online platform. The patient is granted gradual access to content such as video clips, texts, audio files, and work sheets, and the treatment is guided and supported by a clinician through an embedded asynchronous message system. The basic principle is that internet-delivered treatment should target the same core treatment processes as face-to-face treatment, with only the mode of delivery changing [12]. Internet-delivered treatment compares favorably with face-to-face treatment as it is unrestrained by geographical distance and interferes less with work and family obligations. Thus, it may inflict less strain on the patients' daily activities. Furthermore, studies suggest that internet-delivered treatment is less stigmatizing and therefore less of a barrier for seeking treatment than face-to-face treatment [13]. In addition, compared to treatment in an outpatient clinic, internet-delivered treatment may involve fewer societal costs due to absence from work and travel expenses.

Internet-delivered treatment for common mental disorders has been investigated in well over 100 randomized trials, and a recent meta-analysis showed that internet-delivered cognitive-behavioral therapy (iCBT) for depressive and anxiety disorders can be just as effective as face-to-face-treatment [14]. In 3 randomized controlled trials (RCTs), Hedman-Lagerlöf and colleagues showed that iCBT can be highly effective for severe health anxiety [15-17]. Studies have also shown that the treatment effect is long-term and cost-effective [18].

Poor adherence to internet-delivered treatment is a widespread and well-documented concern [19,20], which may limit its potential effectiveness [21]. A recent meta-analysis found that the mean adherence to protocol was higher in internet-delivered acceptance and commitment therapy (iACT) compared to iCBT [22]. Acceptance and commitment therapy (ACT) is a new, promising treatment within the third wave of cognitive-behavioral therapies, which has shown a positive effect across a range of mental health conditions, including health anxiety [23-26]. ACT aims to increase "psychological flexibility" which is considered a core process across different disorders [27]. Patients with severe health anxiety report low psychological flexibility, which often narrows the behavioral responses to health anxiety [28]. Patients spend a lot of time and energy trying to control or avoid thoughts, feelings, and sensations associated with illness, limiting the focus on value-based activities and creation of a meaningful life. ACT deliberately does not focus on symptom reduction, since this focus can maintain the very same symptoms by continuing to control or avoid them. Instead ACT aims to improve 2 skills: acceptance of inner experiences and commitment to values-based behaviors. A recent trial on iACT for social anxiety and panic disorder suggested that iACT may be feasible for treating anxiety disorders [29]. However, to our knowledge, iACT for severe health anxiety has not previously been developed or tested.

The aims of this study were to (1) describe the development of a new treatment concept for patients with severe health anxiety encompassing patient self-referral, video-based diagnostic assessment, and treatment with iACT over 12 weeks; and (2) evaluate the feasibility and overall outcome of this new treatment concept. The feasibility evaluation included patients' feedback provided both at the video-based assessment and post-treatment interview, self-reported questionnaires, and the patients' attrition and adherence to the treatment. The evaluation did not apply a priori defined criteria of success, since this study was a proof of concept prior to conducting a larger, RCT.

Methods

Phase I: Development

Both the technical platform and the contents of the treatment were developed de novo (Multimedia Appendix 1). The treatment content was developed according to an evidence-based manual [30]. Decisions not informed by previous research were made in collaboration with an expert group consisting of 2 psychologists and 1 psychiatrist from the Research Clinic for Functional Disorders and Psychosomatics at Aarhus University Hospital in Denmark. This group was supplemented with 2 external psychologists with expertise in developing and conducting research on internet-delivered programs. Two focus group interviews with former patients were conducted to establish relevant themes to address in new treatment content (eg, patient videos). The technical platform was developed in collaboration with a Web-developer. New procedures—patient self-referral and video-based assessment—were applied.

Self-Referral and Video-Based Assessment

Self-referral was developed and tested as a new treatment entry through the clinic's webpage, where patients could log into a portal using their unique Danish personal identification number. The self-referral procedure consisted of the following steps: (1) patients gave written consent for the clinician to access information on their health status through the electronic patient record prior to assessment, (2) patients were asked to describe their problem in an open text field "Please describe your health anxiety in your own words and how you feel at the moment," and (3) patients completed baseline screening questionnaires measuring health anxiety symptoms among others. Video-based assessment was implemented to support nationwide recruitment. A shortened version of the semi-structured psychiatric interview, Schedules for Clinical Assessment in Neuropsychiatry (SCAN) [31], was adapted for this purpose in order to focus and shorten the assessment. The final version was screened for severe health anxiety and the corresponding diagnoses of hypochondriasis, illness anxiety disorder, and somatic symptom disorder. Furthermore, depression, anxiety disorders including obsessive-compulsive disorder and other somatoform disorders according to the International Classification of Diseases (ICD-10) were assessed.

Technical Platform

Accessibility, interactivity, contact, and technical solution are some of the focal points that define the functionality and appearance of internet-delivered treatment programs. The decisions regarding these focal points as they relate to the present study are described below.

Accessibility

Accessibility can range from full, open access to health care integrated programs with clinician-administered inclusion. Since this platform was developed and operated in a hospital setting, access was administered by health professionals to secure that the hospital's legal responsibility was ensured. Treatment progress was monitored and access to treatment modules was granted consecutively by the clinician.

Interactivity

Interactivity can range from fixed content with no interaction to more responsive content such as online exercises or games that are adapted based on the patient's interaction. The platform's content was designed to be relatively fixed (text, audio files, video clips, illustrations), but included interactive work sheets which were completed by the individual patient using embedded text fields. All sheets were automatically collected and stored in a folder on the front page and shared with the patient's personal clinician.

Contact

Contact can range from pure self-help to varying degrees and formats of contact (eg, written/telephone/video support to blended care where online tools are blended with face-to-face contact). The current platform enabled written, asynchronous clinician guidance (no real-time chat). This format was chosen to minimize interference with the patient's daily activities. Guidance was provided throughout the 12 weeks of treatment. Every patient had a designated personal clinician and messages were answered within 48 hours on weekdays. The content of the written guidance was not predetermined or restricted, and no limit was established concerning contact frequency and length.

Technical Solution

Technical solution can range from tools designed for mobile apps to extensive Web-based programs. The current platform was developed as a Web-based app built upon Drupal, a flexible open source content management system (CMS) and development framework. This solution was chosen because the treatment material is comprehensive and specialized functionality had to be added to the platform. The following solutions were developed: (1) a 2-factor authentication login which interfaces with the Danish public login system (NemID); (2) an embedded, securely encrypted message system for written communication; (3) an automatic text message (short message system, SMS) notification system (eg, notifies patients about new content such as messages, modules, questionnaires); and (4) a clinician-monitored control panel with automatic notifications with information about patient activity (eg, new messages, work sheets, activity, or missed questionnaires). In addition, the responsive design allows access to the treatment program through mobile devices and tablets.

Treatment Content: Intervention

The treatment content was based on an empirically tested group-based ACT program for patients with severe health anxiety [24,30], and clinical experience with the manual and focus group interviews with former patients receiving group treatment. Literature on internet-delivered treatments was also reviewed. All treatment content aims to target and affect core structures of health anxiety such as experiential avoidance of unpleasant illness-related thoughts, feelings, and bodily sensations that inhibit values-based actions. The treatment program trains acceptance of these unpleasant inner states, clarifies values, and simultaneously promotes values-based exposure. The overall intention is to improve the patient's psychological flexibility (ie, ensure they are able to experience

a more flexible and value-based behavioral repertoire when health anxiety is present). This includes the distinction between what can and cannot be changed. Since ACT is a behavior therapy, the treatment works through both the processes of accepting what cannot be controlled (thoughts, feelings, sensations), and change what can be controlled (behaviors). The progression of treatment processes is illustrated in [Table 1](#).

The number of modules in the original face-to-face manual was shortened from 10 to 7 ([Textbox 1](#)). Since this program was primarily based on self-help, the language was revised and simplified so that all content elements were self-explanatory. The front page features 10 widgets (messages, work sheets, acute help, and modules 1 to 7) ([Multimedia Appendices 2 and 3](#)). Each module has between 10 to 15 pages of content which is primarily text-based. The modules also contain audio files with guided exercises, such as mindfulness training, and patient videos with 5 former patients sharing their experience of health anxiety and treatment processes. The videos were produced to enhance treatment comprehension, recognition, and patient motivation. All videos and audio exercises are relatively short (between 4 to 20 minutes), and each module introduces at least 1 new video and audio exercise. Furthermore, illustrations were hand drawn, graphic design was added to structure different types of treatment elements, and written work sheets were embedded. All modules follow the same structure; the first page is an introduction to the following theme and a repetition of the previous one, next follows the main treatment content, and lastly a summary of the current module, clarifying questions and an introduction to weekly homework exercises. The homework consists of 2 alternating exercises each week: an audio exercise primarily based on mindfulness and a behavioral exercise based on values-based exposure ([Multimedia Appendices 4-6](#)).

Phase II: Design of Feasibility Testing

Study Design and Setting

This feasibility study took place from April 2015 to October 2015 at the Research Clinic for Functional Disorders and Psychosomatics at Aarhus University Hospital in Denmark. It was an open trial with no control group. Self-report questionnaires were obtained at baseline prior to diagnostic assessment, at post-treatment, and at 3-month follow-up. Patients

not completing any modules were excluded from the final analyses as the primary aim of the study was to evaluate the feasibility of the new treatment set-up and program. Treatment completion was defined a priori as 3 or more modules completed since the first 2 modules primarily consisted of psycho-education and did not explicitly include behavior change strategies. Semi-structured interviews with patients after end of treatment were conducted by the clinician assigned to the individual patient to explore the usability and acceptability of the procedure with self-referral, video-based assessment, as well as the treatment program. All participants provided written informed consent, and the study was approved by the Danish Data Protection Agency, Central Denmark Region (ID no. 1-16-02-427-14).

Participant Recruitment and Eligibility

A total of 15 patients were recruited, 10% of the projected sample size for the RCT. This was considered sufficient for the feasibility evaluation of the treatment concept including self-referral, assessment, and treatment. Patients were recruited through the clinic's webpage, which announced information about the study, and presented a patient video about health anxiety. Patient self-referrals were screened for eligibility by a psychologist (first author) and then allocated to a video-based clinical assessment. The assessments were conducted by psychologists or psychiatrists trained in the use of the shortened version of SCAN [31]. Health information was assessed and the electronic patient record was examined before the interview. A psychiatrist provided supervision in regard to somatic questions. Eligible participants had to (1) report symptoms corresponding to fulfillment of the empirically-based diagnostic criteria for severe health anxiety in accordance with Fink et al [1]; (2) have a Whiteley Index-7 (WI-7) score above 21.4 (scale range 0 to 100), which is established as a clinically relevant cut-off score [10,24]; (3) be at least 18 years old; (4) be able to read and understand Danish; and (5) have a primary diagnosis of health anxiety if co-morbid disorders were present. Patients were excluded from the study if they (1) were at risk for suicide; (2) had current or previous episodes of psychosis; (3) had current abuse of alcohol, drugs, or medication; (4) were pregnant; and (5) did not give informed consent to the study.

Table 1. Treatment processes.

Module	Functional analysis	Mindfulness	Willingness and commitment	Values	Defusion	Acceptance
1	✓			✓		
2	✓	✓		✓		
3		✓	✓			✓
4		✓	✓	✓		
5		✓	✓	✓	✓	✓
6		✓		✓	✓	✓
7	✓					✓

Textbox 1. Overview of the treatment.

Treatment targets for each module

Module 1: The first important step

- What are health anxiety and acceptance and commitment therapy?
- Overview and introduction to the treatment
- Motivation and values for participation

Module 2: Your experiences managing health anxiety

- Model of health anxiety and introduction to values
- Control, avoidance and creative hopelessness
- Mindfulness and rumination about illness

Module 3: Challenging and accepting anxiety

- Evolutionary function of bodily symptoms of anxiety
- The workability of control and avoidance behavior
- Challenging (exposure) and accepting anxiety

Module 4: You get to know your values when taking steps

- Values sorting
- Values compass
- Values-based exposure-hierarchy

Module 5: What is stopping me?

- Inner barriers (thoughts, feelings, bodily sensations)
- Defusion and acceptance
- Mindfulness and self-as-context

Module 6: Self-compassion and boundaries

- How stress affects bodily symptoms
- Balancing maladaptive “all-or-nothing” behavior
- Personal boundaries, open up and self-care

Module 7: Relapse prevention

- Repetition of treatment principles
- Consolidating personal insights from the treatment
- Normalization of relapse and relapse prevention

Measures

The mean sum score of the scales presented below were transformed into a 0 to 100 score point scale according to equation 1:

$$(1) \text{ Scale} = [(score - minimum) \div (maximum - minimum)] \times 100$$

This was chosen to facilitate the comparison of changes between measures in this and previous studies [24,32].

Primary Outcome

The WI-7 [33] is a 7-item measure of health anxiety symptoms including items such as “Do you often worry about the possibility that you have a serious illness?” Each item is scored

on a 5-point Likert scale ranging from 0 to 4 (range 0 to 28), with higher scores indicating more severe health anxiety. WI-7 is broadly employed and has shown acceptable psychometric properties in primary care [10], and good sensitivity and specificity in screening for DSM-IV hypochondriasis [34].

Secondary Outcomes

The Symptom Checklist-92 (SCL-92) [35] subscale for depression (SCL-92 Dep) is a 13-item measure of depression (range 13 to 65). The subscale for anxiety (SCL-92 Anx) is a 10-item measure of anxiety (range 10 to 50). Each item is scored on a 5-point Likert scale ranging from 1 to 5, with higher scores indicating more severe illness.

The Short-Form Health Survey (SF-12) [36,37] with 12 items measures 2 dimensions of health-related quality of life (HRQoL): the physical component summary (PCS) and the mental component summary (MCS). Both scales range from 0 to 100, with higher scores indicating better physical and mental HRQoL.

Life satisfaction [38] is a 1-item measure scored on a 10-point Likert scale ranging from 0 to 10, where 0 and 10 encompass the “the worst possible life” and “the best possible life,” respectively.

The Acceptance and Action Questionnaire-II (AAQ-II) [39,40] is a 7-item measure of psychological flexibility. Each item is scored on a 7-point Likert scale ranging from 1 to 7 (range 10 to 70), with higher scores indicating better functioning.

Statistical Analysis

Data were summarized using means and standard deviations. The mean differences from baseline to post-treatment and from baseline to 3-month follow-up were analyzed using paired *t* tests. The assumptions behind the paired *t* test were assessed by graphical inspection of a quantile-quantile (Q-Q) plot of the differences and a Bland-Altman plot. An estimate of within group effect sizes was calculated using the standardized response mean (SRM) according to equation 2, by the formula, where X_1 and X_2 are the means of the measurements at time points 1 and 2, respectively. SD_{DIFF} is the standard deviation of the variable containing the difference between the 2 measurements.

$$(2) \text{ SRM} = (X_1 - X_2) \div SD_{DIFF}$$

Adherence was measured as the mean number of modules completed, and exchanged messages in the program were summarized using means and standard deviations. All analyses were performed using Stata version 13 for Windows. The post-treatment interviews were conducted using semi-structured questions. The answers were not analyzed quantitatively.

Results

Patient Characteristics and Adherence

A total of 18 patients were self-referred of which 3 (3/18, 17%) withdrew prior to assessment (Figure 1). Of the patients, 15 (15/18, 83%), including 12 females (12/18, 67%) were assessed for eligibility, all of whom met the inclusion criteria and were included in the study. Socio-demographic and clinical details are shown in Table 2. Of the patients, 80% (12/15) completed treatment (ie, performed a minimum of 3 modules). The majority of the completers were considered active during all 12 weeks of treatment, independently of the number of modules finalized. One patient (1/15, 7%) dropped out prior to module 1 due to technical obstacles. Thus, post-treatment data on 14 (14/15, 93%) patients and 3-month follow-up data on 12 (12/15, 80%) patients were included in the final statistical analyses (Figure 1).

Of the patients, 12 (12/15, 80%) agreed to participate in the semi-structured interviews at post-treatment. Due to

cancellations, only 10 interviews were conducted. The patient who dropped out and 1 of the non-completers participated in the interview ensuring a varied evaluation.

Feasibility of Patient Recruitment and Assessment

Online patient recruitment through the clinic’s webpage attracted the patient population of interest. All self-referred patients were judged to have health anxiety as their primary problem based on the first screening and they were invited for assessment. Patients reported that it was easy to find the trial and go through the steps of self-referral.

At the end of the video-based interview, patients commented on their user experience. Overall, patients found it convenient, and the majority expressed feeling more secure at home than having to go to the hospital. A few patients felt anxious prior to the assessment due to the new video-format, and technical issues did interrupt a few interviews. In spite of this, most patients expressed that the video-format did not interfere negatively with their sense of contact with the clinician during the assessment. All clinicians had no prior experience with the video-format, and they reported positively about the amount of clinical information the assessment allowed for.

Feasibility of the Treatment Platform and Content

The post-treatment patient interviews revealed overall satisfaction with relevance and progression of treatment themes and a positive response to the use of patient videos. There was no consistent pattern in content feedback (eg, some patients emphasized the importance of mindfulness exercises, and some patients did not use them at all). Patients sent an average of 12.9 (SD 7.76) messages (range 4-29) to their clinician, and clinicians sent an average of 18.5 (SD 3.46) messages (range 12-25) to their patient. Some patients expressed that this form of communication allowed for more honesty compared with earlier experiences with face-to-face therapy, whereas for others it was a hindrance. Therefore, some patients wrote many extensive messages where others consulted the clinician only to a limited extent. The clinicians reported that it was possible to have a good clinical sense of the patient. However, for those patients who did not use the clinician, the therapeutic reinforcement was missing to a larger extent. Furthermore, the patients were generally surprised by the amount of time they needed to complete a module.

Changes in Outcome Measures

Mean, standard deviation, and percent changes at baseline, post-treatment, and 3-month follow-up are shown in the Table 3. Mean difference, CI, *P* values, and effect sizes are shown in Table 4. Patients reported statistically significant reductions in health anxiety symptoms from baseline to post-treatment with a difference in WI-7 score of 37.2 points (95% CI 24.1-50.4, $t_{13} = 6.11$, $P < .001$), and at 3-month follow-up with a difference of 33.9 points (95% CI 13.6-54.3, $t_{11} = 3.66$, $P = .004$) with large within-group effect sizes (SRM 1.06-1.63).

Figure 1. Study flowchart.




Table 2. Baseline characteristics of study sample (N=15).

Variable	Value
Age (years), mean (range)	38.8 (20.8-56.6)
Sex, n (%)	
Female	12 (80)
Male	3 (20)
Education, n (%)	
Basic school (9-15 years)	9 (60)
Further/higher education (<4 years)	6 (40)
Work status, n (%)	
Full-time employed	7 (47)
Part-time employed	3 (20)
Unemployed	0 (0)
Disability pension or flexible work	4 (27)
Other (eg, maternity leave)	1 (7)
Absence from work, n (%)	
No absence	10 (67)
Full-time absence	3 (20)
Part-time absence	2 (13)
Living status, n (%)	
Alone	2 (13)
With someone	13 (87)
Duration of health anxiety (years), mean (range)	15.2 (0.7-30)
Onset of health anxiety (years), mean (range)	23.6 (9-40)
Psychiatric co-morbidity, n (%)	
Other anxiety disorders	4 (27)
Depression	0 (0)
Other somatoform disorders ^a	2 (13)

^aSomatization disorder or undifferentiated somatoform disorder.

Table 3. Mean, standard deviation and percent change at baseline, post-treatment, and 3-month follow-up. WI-7: Whiteley-7 Index. SCL-92 Dep: symptom checklist-92 depression. SCL-92 Anx: symptom checklist-92 anxiety. PCS: physical component summary. SF-12: short-form health survey. MCS: mental component summary. AAQ: acceptance and action questionnaire.

Outcome measure ^a	Baseline (T1; N=15)		Post-treatment (T2; N=14)		3-month follow-up (T3; N=12)	
	Mean (SD)		Mean (SD)	% ^b	Mean (SD)	% ^c
Primary outcome						
Health anxiety (WI-7)	80.9 (18.5)		43.6 (13.0)	-46	45.5 (29.6)	-44
Secondary outcomes						
Depression (SCL-92 Dep)	40.4 (13.2)		21.2 (13.3)	-48	16.2 (13.6)	-60
Anxiety (SCL-92 Anx)	43.8 (14.3)		27.5 (16.4)	-37	18.8 (12.6)	-57
PCS (SF-12)	44.7 (15.8)		46.2 (11.8)	3	48.5 (13.8)	9
MCS (SF-12)	28.7 (9.5)		43.6 (10.0)	52	44.3 (12.1)	54
Life satisfaction	45.7 (12.2)		59.3 (18.2)	30	71.7 (14.0)	57
Process measure						
Psychological flexibility (AAQ-II)	37.9 (15.8)		61.4 (20.0)	62	69.6 (16.3)	84

^aHigh score is many symptoms, except for life satisfaction, SF-12, and psychological flexibility. Scale 0 to 100.

^bPercent change from baseline to post-treatment, measured by $((T1-T2)/T1) \times 100$.

^cPercent change from baseline to 3-month follow-up, measured by $((T1-T3)/T1) \times 100$.

Table 4. Change in outcome measures and effect sizes. MD: mean difference. SRM: standardized response mean. WI-7: Whiteley-7 Index. SCL-92 Dep: symptom checklist-92 depression. SCL-92 Anx: symptom checklist-92 anxiety. PCS: physical component summary. SF-12: short-form health survey. MCS: mental component summary. AAQ: acceptance and action questionnaire.

Outcome measure	Baseline to post-treatment (N=14)				Baseline to 3-month follow-up (N=12)			
	MD	95% CI	P value	SRM	MD	95% CI	P value	SRM
Primary outcome								
Health anxiety (WI-7)	37.2	24.1 to 50.4	<.001	1.63	33.9	13.6 to 54.3	.004	1.06
Secondary outcomes								
Depression (SCL-92 Dep)	19.2	11.2 to 27.3	<.001	1.38	21.5	10.5 to 32.4	.001	1.24
Anxiety (SCL-92 Anx)	16.3	3.3 to 29.2	.02	0.73	25.6	14.9 to 36.4	<.001	1.51
PCS (SF-12)	-3.3	-11.9 to 5.3	.42	0.23	-5.1	-13.3 to 3.0	.19	0.42
MCS (SF-12)	-15.0	-21.0 to -9.0	<.001	1.50	-15.3	-21.7 to -8.9	<.001	1.61
Life satisfaction	-13.6	-23.9 to -3.3	.01	0.76	-27.5	-37.3 to -17.7	<.001	1.78
Process measure								
Psychological flexibility (AAQ-II)	-23.5	-35.3 to -11.7	<.001	1.15	-31.2	-42.1 to -20.2	<.001	1.80

There were no significant changes in scores from post-treatment to 3-month follow-up which suggests stability of clinical outcome. Furthermore, patients reported significant reductions in symptoms of anxiety and depression, and significant increases in life satisfaction and psychological flexibility (Table 4). Effect sizes were moderate to large on all secondary outcomes (SRM 0.73-1.80), with the exception of the SF-12 PCS measuring physical functioning. Apart from life satisfaction, which further increased significantly from post-treatment to 3-month follow-up (SRM 0.96), there were no significant changes in scores from post-treatment to 3-month follow-up.

Subsequent Revisions of the Treatment Program and Set-Up

The patient interviews gave rise to revision and further extension of some treatment themes and exercises. Furthermore, we learned that patients need to not only be motivated for internet-delivered treatment, but also to be able to work independently, prioritize their time with regard to treatment engagement, and have a minimum of technical know-how. To further improve adherence prior to the RCT, a patient recruitment video was developed to provide a clear picture of the treatment format, emphasizing the time and effort needed to participate in this type of treatment. This was also further addressed in the assessment interview. Patients who were ambivalent had up to 2 weeks to decide whether to enter the

randomization for the following trial. Based on the experience with the drop-out of 1 patient due to technical obstacles, we included a question about computer skills in the baseline questionnaire. We also enquired more about this in the assessment interview. Finally, due to discussions raised about changes in medication during the pilot study, an extra inclusion criterion was added for the RCT to ensure that patients had been stable on anxiety medication for at least 2 months prior to inclusion.

Discussion

Principal Findings

The main finding of this pilot feasibility study was that iACT encompassing online self-referral, video-based assessment, and internet-delivered treatment was a feasible treatment concept for patients suffering from severe health anxiety. We found overall positive outcomes with large improvements in health anxiety symptoms and medium to large improvements in anxiety, depression, HRQoL, and life satisfaction, which improved further during the 3-month follow-up period. The observed reductions in health anxiety symptoms were similar to those reported in a previous, uncontrolled, face-to-face pilot study, and a subsequent RCT on group-delivered ACT for patients with health anxiety [24,32]. Since health anxiety is considered a chronic disorder if untreated [10], spontaneous remission is unlikely to have contributed significantly to this effect. It is often debated whether symptom measures should be used as a primary outcome in ACT trials, since the treatment does not focus on symptom reduction but instead aims to increase behavioral flexibility and valued living. In our study, we also measured psychological flexibility, which did show the largest improvement (SRM 1.80). This supports the model of change suggesting that increased psychological flexibility is associated with general psychological well-being and might even precede symptom reduction [28,41]. Thus, symptom reduction is a welcomed treatment outcome but should not be mistaken as a therapeutic goal in treatment. In the present study, only 1 (1/15, 7%) patient dropped out, and 12 patients (12/15, 80%) completed 12 weeks of treatment. This was comparable to iCBT for health anxiety which has shown a similar completion rate of 85% [15]. Since the majority of patients completed the treatment, iACT seems to be an acceptable and feasible treatment for severe health anxiety.

Strengths and Limitations

The thorough standardized clinical assessment and well-defined diagnostic inclusion criteria were strengths of this pilot study. Moreover, both completers and non-completers participated in the post-treatment interview and most data were available for patients at follow-up. Lastly, the study employed validated clinical outcome measures. Even though this study did not employ a randomized design, most outcomes pointed in the same direction: symptoms improved following iACT. The most important limitation of this study was the uncontrolled study design and small sample size. The design was chosen based on the aims of this study, and the sample size did not limit the feasibility evaluation, which was the primary aim. The preliminary observed clinical changes should be replicated in

a larger RCT to establish clinical relevance. Another limitation was that the video-based assessment was not evaluated systematically. We did discuss the technical and clinical implications at supervision, and asked the patients about their experience at the interview, but we did not collect our experiences in a structured manner. Lastly, the post-treatment patient interviews were not analyzed systematically and since they were conducted by the dedicated clinician, there was a risk of bias due to socially desirable answers.

Patient Recruitment and Generalizability

Patient self-referral is rarely applied in a publicly funded health care system based on gatekeeping, where patients are referred primarily through the general practitioner. We found that the majority of those who self-referred fulfilled criteria for severe health anxiety, which may indicate that self-referral is a sensible recruitment procedure in patients with severe health anxiety. One explanation may be that patients with health anxiety often seek medical information via the internet, and may have come across the clinic's webpage while searching for explanations for their symptoms. The patient video on the webpage was likely to have further increased recruitment sensitivity as many patients reported that they recognized their own health anxiety from the patient's stories. Former studies support that patient self-referral is a feasible recruitment method for internet-delivered psychological treatment [42,43]. However, patient characteristics may differ compared to patients referred from the general practitioner to outpatient treatment. Prior to this study, we anticipated that the treatment setup would appeal to males and younger patients with less severe symptoms of health anxiety. Studies tend to find no age or gender difference in the distribution of health anxiety [10], yet males and younger patients seem to be less frequently referred to treatment [24]. However, in the present study, we recruited 80% (12/15) female patients, with an average age of 38.8 years. Moreover, patients had health anxiety for approximately 15 years and were quite impaired at referral, with an average WI-7 score of 81 (range 0 to 100). Online administration of questionnaires has shown to be a valid method for measuring symptoms of health anxiety, which suggests that the surprising level of symptom severity cannot be explained by this [44]. This might imply that a different patient subgroup was reached, even though the group did not differ in the expected direction. In the DSM-5, patients suffering from health anxiety are categorized as either care-seeking or care-avoiding [2]. It may be that self-referral appeals to more avoidant patients with health anxiety, thus making treatment accessible for a different subgroup. However, these speculations need to be investigated in a larger patient sample.

Treatment Processes and Clinical Implications

The overall aim of iACT for health anxiety is to target and improve a patient's psychological flexibility, which previously has shown to be a central process of change [28,45]. This was done through several treatment processes as previously illustrated (Textbox 1). Specifically, we had a distinct focus on acceptance of inner experiences and values in order to broaden the behavioral repertoire from control and avoidance to values-based behaviors. This involved gradually changing

behaviors from unworkable short-term strategies to long-term goals in line with the chosen values. The distinct focus on personal values in ACT compared to CBT may explain why iACT tends to have better adherence to protocol compared to iCBT [22]. Future studies should include process measures to further understand the role of values and investigate the mediational effect on symptoms.

Clinician guidance was added to this treatment program because studies generally find that unguided internet-delivered treatment has higher attrition [20,46]. The asynchronous message system enabled a more flexible and patient-tailored contact in terms of individual needs of guidance. This was also reflected in the wide range of messages exchanged between each patient and their allocated clinician. The aim of the written guidance was to motivate patients, to validate patient efforts, to clarify misunderstandings, and to personalize treatment content. This study did not measure the quality of therapeutic alliance, but a former review on internet-delivered treatment across patient populations found client-rated alliances close to those found in face-to-face studies [47]. However, the review also reported

that clinicians rated the alliance lower than in face-to-face treatment. This is in line with the feedback from clinicians in the present study regarding patients who did not use their dedicated clinician. This may suggest that clinicians are affected more by the internet-delivered treatment modality than patients.

Conclusion

Severe health anxiety is a disabling and costly disorder with a lack of accessible, specialized treatment options. Here, we showed that online self-referral, video-based assessment, and iACT may be a feasible treatment set-up for patients suffering from severe health anxiety. The promising outcome results also suggest potential efficacy for this iACT program which is currently being tested in a larger RCT. If proven effective, iACT may increase availability and accessibility of specialized treatment for health anxiety. Future studies need to evaluate the sensitivity of patient self-referral, and to compare characteristics in patients self-referred or referred through a health care professional, to determine whether different patient populations are reached.

Acknowledgments

We thank our colleague, Web developer Nicolaj Knudsen from the Research Clinic for Functional Disorders and Psychosomatics at Aarhus University Hospital, for his expertise and patience when developing the platform for internet-delivered treatment. We would also like to thank Jens S Jensen for statistical assistance and Morten Pilegaard for language revision. We would also like to show our gratitude to Susanne Daugaard and Britta Ravn from the Center for Telemedicine in Central Region Denmark, for their support and guidance. This work was supported by the TRYG foundation (ID no. 102644). The funding sources were not involved in designing, conducting, or preparing this manuscript for publication.

Conflicts of Interest

EHL and BL are shareholders of DahliaQomit Inc, a company specializing in online psychiatric symptom assessment.

Multimedia Appendix 1

Video demonstration of the program.

[[MP4 File \(MP4 Video\), 10MB - mental_v5i2e28_app1.mp4](#)]

Multimedia Appendix 2

iACT program.

[[PNG File, 934KB - mental_v5i2e28_app2.png](#)]

Multimedia Appendix 3

Graphic design elements aiming to structure the treatment content throughout the program.

[[PNG File, 46KB - mental_v5i2e28_app3.png](#)]

Multimedia Appendix 4

Illustration of long term consequences of health anxiety.

[[PNG File, 7MB - mental_v5i2e28_app4.png](#)]

Multimedia Appendix 5

Illustration of control and avoidance as a mean to escape unpleasant thoughts, feelings and bodily sensations; here illustrated as 3 monsters.

[[PNG File, 4MB - mental_v5i2e28_app5.png](#)]

Multimedia Appendix 6

Illustration of the principle of graded exposure.

[[PNG File, 283KB - mental_v5i2e28_app6.png](#)]

References

1. Fink P, Ørnbøl E, Toft T, Sparle KC, Frostholt L, Olesen F. A new, empirically established hypochondriasis diagnosis. *Am J Psychiatry* 2004 Sep;161(9):1680-1691. [doi: [10.1176/appi.ajp.161.9.1680](#)] [Medline: [15337660](#)]
2. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders, 5th Edition*. Arlington, VA: American Psychiatric Association; 2013.
3. Creed F, Barsky A. A systematic review of the epidemiology of somatisation disorder and hypochondriasis. *J Psychosom Res* 2004 Apr;56(4):391-408. [doi: [10.1016/S0022-3999\(03\)00622-6](#)] [Medline: [15094023](#)]
4. Tyrer P, Cooper S, Crawford M, Dupont S, Green J, Murphy D, et al. Prevalence of health anxiety problems in medical clinics. *J Psychosom Res* 2011 Dec;71(6):392-394. [doi: [10.1016/j.jpsychores.2011.07.004](#)] [Medline: [22118381](#)]
5. Weck F, Richtberg S, Neng J. Epidemiology of hypochondriasis and health anxiety: comparison of different diagnostic criteria. *CPSR* 2014 Jan 31;10(1):14-23. [doi: [10.2174/1573400509666131119004444](#)]
6. Sunderland M, Newby JM, Andrews G. Health anxiety in Australia: prevalence, comorbidity, disability and service use. *Br J Psychiatry* 2013 Jan;202(1):56-61 [FREE Full text] [doi: [10.1192/bjp.bp.111.103960](#)] [Medline: [22500013](#)]
7. Thomson AB, Page LA. Psychotherapies for hypochondriasis. *Cochrane Database Syst Rev* 2007 Oct 17(4):CD006520. [doi: [10.1002/14651858.CD006520.pub2](#)] [Medline: [17943915](#)]
8. Tyrer P, Eilenberg T, Fink P, Hedman E, Tyrer H. Health anxiety: the silent, disabling epidemic. *BMJ* 2016 Dec 25;353:i2250. [Medline: [27112356](#)]
9. Barsky AJ, Ettner SL, Horsky J, Bates DW. Resource utilization of patients with hypochondriacal health anxiety and somatization. *Med Care* 2001 Jul;39(7):705-715. [Medline: [11458135](#)]
10. Fink P, Ørnbøl E, Christensen KS. The outcome of health anxiety in primary care. A two-year follow-up study on health care costs and self-rated health. *PLoS One* 2010 Mar 24;5(3):e9873 [FREE Full text] [doi: [10.1371/journal.pone.0009873](#)] [Medline: [20352043](#)]
11. Mykletun A, Heradstveit O, Eriksen K, Glozier N, Øverland S, Maeland JG, et al. Health anxiety and disability pension award: the HUSK study. *Psychosom Med* 2009 Apr;71(3):353-360. [doi: [10.1097/PSY.0b013e31819cc772](#)] [Medline: [19321853](#)]
12. Hedman E, Ljótsson B, Lindefors N. Cognitive behavior therapy via the Internet: a systematic review of applications, clinical efficacy and cost-effectiveness. *Expert Rev Pharmacoecon Outcomes Res* 2012 Dec;12(6):745-764. [doi: [10.1586/erp.12.67](#)] [Medline: [23252357](#)]
13. Marques L, LeBlanc NJ, Weingarden HM, Timpano KR, Jenike M, Wilhelm S. Barriers to treatment and service utilization in an internet sample of individuals with obsessive-compulsive symptoms. *Depress Anxiety* 2010 May;27(5):470-475. [doi: [10.1002/da.20694](#)] [Medline: [20455248](#)]
14. Andersson G, Cuijpers P, Carlbring P, Riper H, Hedman E. Guided Internet-based vs. face-to-face cognitive behavior therapy for psychiatric and somatic disorders: a systematic review and meta-analysis. *World Psychiatry* 2014 Oct;13(3):288-295 [FREE Full text] [doi: [10.1002/wps.20151](#)] [Medline: [25273302](#)]
15. Hedman E, Andersson G, Andersson E, Ljótsson B, Rück C, Asmundson GJ, et al. Internet-based cognitive-behavioural therapy for severe health anxiety: randomised controlled trial. *Br J Psychiatry* 2011 Mar;198(3):230-236 [FREE Full text] [doi: [10.1192/bjp.bp.110.086843](#)] [Medline: [21357882](#)]
16. Hedman E, Axelsson E, Göring A, Ritzman C, Ronnheden M, El AS, et al. Internet-delivered exposure-based cognitive-behavioural therapy and behavioural stress management for severe health anxiety: randomised controlled trial. *Br J Psychiatry* 2014 Oct;205(4):307-314 [FREE Full text] [doi: [10.1192/bjp.bp.113.140913](#)] [Medline: [25104835](#)]
17. Hedman E, Axelsson E, Andersson E, Lekander M, Ljótsson B. Exposure-based cognitive-behavioural therapy via the internet and as bibliotherapy for somatic symptom disorder and illness anxiety disorder: randomised controlled trial. *Br J Psychiatry* 2016 Dec;209(5):407-413 [FREE Full text] [doi: [10.1192/bjp.bp.116.181396](#)] [Medline: [27491531](#)]
18. Hedman E, Andersson E, Lindefors N, Andersson G, Rück C, Ljótsson B. Cost-effectiveness and long-term effectiveness of internet-based cognitive behaviour therapy for severe health anxiety. *Psychol Med* 2013 Feb;43(2):363-374. [doi: [10.1017/S0033291712001079](#)] [Medline: [22608115](#)]
19. Kelders SM, Kok RN, Ossebaard HC, Van Gemert-Pijnen JE. Persuasive system design does matter: a systematic review of adherence to web-based interventions. *J Med Internet Res* 2012 Nov 14;14(6):e152 [FREE Full text] [doi: [10.2196/jmir.2104](#)] [Medline: [23151820](#)]
20. Christensen H, Griffiths KM, Farrer L. Adherence in internet interventions for anxiety and depression. *J Med Internet Res* 2009 Apr 24;11(2):e13 [FREE Full text] [doi: [10.2196/jmir.1194](#)] [Medline: [19403466](#)]

21. Donkin L, Christensen H, Naismith SL, Neal B, Hickie IB, Glozier N. A systematic review of the impact of adherence on the effectiveness of e-therapies. *J Med Internet Res* 2011 Aug 05;13(3):e52 [FREE Full text] [doi: [10.2196/jmir.1772](https://doi.org/10.2196/jmir.1772)] [Medline: [21821503](https://pubmed.ncbi.nlm.nih.gov/21821503/)]
22. Brown M, Glendenning A, Hoon AE, John A. Effectiveness of Web-delivered acceptance and commitment therapy in relation to mental health and well-being: a systematic review and meta-analysis. *J Med Internet Res* 2016 Aug 24;18(8):e221 [FREE Full text] [doi: [10.2196/jmir.6200](https://doi.org/10.2196/jmir.6200)] [Medline: [27558740](https://pubmed.ncbi.nlm.nih.gov/27558740/)]
23. Hacker T, Stone P, MacBeth A. Acceptance and commitment therapy - do we know enough? Cumulative and sequential meta-analyses of randomized controlled trials. *J Affect Disord* 2016 Jan 15;190:551-565. [doi: [10.1016/j.jad.2015.10.053](https://doi.org/10.1016/j.jad.2015.10.053)] [Medline: [26571105](https://pubmed.ncbi.nlm.nih.gov/26571105/)]
24. Eilenberg T, Fink P, Jensen JS, Rief W, Frosthalm L. Acceptance and commitment group therapy (ACT-G) for health anxiety: a randomized controlled trial. *Psychol Med* 2016 Jan;46(1):103-115. [doi: [10.1017/S0033291715001579](https://doi.org/10.1017/S0033291715001579)] [Medline: [26281857](https://pubmed.ncbi.nlm.nih.gov/26281857/)]
25. A-Tjak JG, Davis ML, Morina N, Powers MB, Smits JA, Emmelkamp PM. A meta-analysis of the efficacy of acceptance and commitment therapy for clinically relevant mental and physical health problems. *Psychother Psychosom* 2015;84(1):30-36. [doi: [10.1159/000365764](https://doi.org/10.1159/000365764)] [Medline: [25547522](https://pubmed.ncbi.nlm.nih.gov/25547522/)]
26. Bluett EJ, Homan KJ, Morrison KL, Levin ME, Twohig MP. Acceptance and commitment therapy for anxiety and OCD spectrum disorders: an empirical review. *J Anxiety Disord* 2014 Aug;28(6):612-624. [doi: [10.1016/j.janxdis.2014.06.008](https://doi.org/10.1016/j.janxdis.2014.06.008)] [Medline: [25041735](https://pubmed.ncbi.nlm.nih.gov/25041735/)]
27. Kashdan TB, Rottenberg J. Psychological flexibility as a fundamental aspect of health. *Clin Psychol Rev* 2010 Nov;30(7):865-878 [FREE Full text] [doi: [10.1016/j.cpr.2010.03.001](https://doi.org/10.1016/j.cpr.2010.03.001)] [Medline: [21151705](https://pubmed.ncbi.nlm.nih.gov/21151705/)]
28. Eilenberg T, Hoffmann D, Jensen JS, Frosthalm L. Intervening variables in group-based acceptance & commitment therapy for severe health anxiety. *Behav Res Ther* 2017 Dec;92:24-31. [doi: [10.1016/j.brat.2017.01.009](https://doi.org/10.1016/j.brat.2017.01.009)] [Medline: [28196772](https://pubmed.ncbi.nlm.nih.gov/28196772/)]
29. Ivanova E, Lindner P, Ly KH, Dahlin M, Vernmark K, Andersson G, et al. Guided and unguided acceptance and commitment therapy for social anxiety disorder and/or panic disorder provided via the Internet and a smartphone application: a randomized controlled trial. *J Anxiety Disord* 2016 Dec;44:27-35. [doi: [10.1016/j.janxdis.2016.09.012](https://doi.org/10.1016/j.janxdis.2016.09.012)] [Medline: [27721123](https://pubmed.ncbi.nlm.nih.gov/27721123/)]
30. Eilenberg T, Frosthalm L, Kronstrand L. Acceptance and commitment group therapy (ACT-G) for severe health anxiety. Aarhus: The Research Clinic for Functional Disorders and Psychosomatics; 2014. URL: http://funktionellelidelser.dk/fileadmin/www.funktionellelidelser.au.dk/Publikationer/ACT_Manual.pdf [accessed 2018-03-27] [WebCite Cache ID [6yDy1jP0d](https://www.webcitation.org/6yDy1jP0d)]
31. World Health Organization. Schedules for Clinical Assessment in Neuropsychiatry. Version 2.1. Geneva: World Health Organization, Division of Mental Health; 1998.
32. Eilenberg T, Kronstrand L, Fink P, Frosthalm L. Acceptance and commitment group therapy for health anxiety--results from a pilot study. *J Anxiety Disord* 2013 Jun;27(5):461-468. [doi: [10.1016/j.janxdis.2013.06.001](https://doi.org/10.1016/j.janxdis.2013.06.001)] [Medline: [23871841](https://pubmed.ncbi.nlm.nih.gov/23871841/)]
33. Fink P, Ewald H, Jensen J, Sørensen L, Engberg M, Holm M, et al. Screening for somatization and hypochondriasis in primary care and neurological in-patients: a seven-item scale for hypochondriasis and somatization. *J Psychosom Res* 1999 Mar;46(3):261-273. [Medline: [10193917](https://pubmed.ncbi.nlm.nih.gov/10193917/)]
34. Conrard M, Cavanagh M, Franklin J, Rief W. Dimensionality of the Whiteley Index: assessment of hypochondriasis in an Australian sample of primary care patients. *J Psychosom Res* 2006 Feb;60(2):137-143. [doi: [10.1016/j.jpsychores.2005.07.003](https://doi.org/10.1016/j.jpsychores.2005.07.003)] [Medline: [16439266](https://pubmed.ncbi.nlm.nih.gov/16439266/)]
35. Derogatis L, Cleary P. Confirmation of the dimensional structure of the SCL-90: a study in construct validation. *J Clin Psychol* 1977 Oct;33(4):981-989. [doi: [10.1002/1097-4679\(197710\)33:4<981::AID-JCLP2270330412>3.0.CO;2-0](https://doi.org/10.1002/1097-4679(197710)33:4<981::AID-JCLP2270330412>3.0.CO;2-0)]
36. Bjorner JB, Damsgaard MT, Watt T, Groenvold M. Tests of data quality, scaling assumptions, and reliability of the Danish SF-36. *J Clin Epidemiol* 1998 Nov;51(11):1001-1011. [Medline: [9817118](https://pubmed.ncbi.nlm.nih.gov/9817118/)]
37. Ware JE. SF-36 health survey update. *Spine (Phila Pa 1976)* 2000 Dec 15;25(24):3130-3139. [Medline: [11124729](https://pubmed.ncbi.nlm.nih.gov/11124729/)]
38. Bendtsen P, Mikkelsen S, Tolstrup J. Youth Profile 2014: Health Behaviour, Health and Well-Being Among Students Under Further Education. Copenhagen: National Institute of Public Health; 2015:1-62.
39. Hayes S, Strosahl K, Wilson K, Bissett R, Pistorello J, Toarmino D, et al. Measuring experiential avoidance: a preliminary test of a working model. *Psychol Rec* 2004 Dec;54(4):553-578 [FREE Full text]
40. Bond F, Hayes S, Baer R, Carpenter KM, Guenole N, Orcutt HK, et al. Preliminary psychometric properties of the Acceptance and Action Questionnaire-II: a revised measure of psychological inflexibility and experiential avoidance. *Behav Ther* 2011 Dec;42(4):676-688. [doi: [10.1016/j.beth.2011.03.007](https://doi.org/10.1016/j.beth.2011.03.007)] [Medline: [22035996](https://pubmed.ncbi.nlm.nih.gov/22035996/)]
41. Hayes SC, Luoma JB, Bond FW, Masuda A, Lillis J. Acceptance and commitment therapy: model, processes and outcomes. *Behav Res Ther* 2006 Jan;44(1):1-25. [doi: [10.1016/j.brat.2005.06.006](https://doi.org/10.1016/j.brat.2005.06.006)] [Medline: [16300724](https://pubmed.ncbi.nlm.nih.gov/16300724/)]
42. Andersson G, Titov N. Advantages and limitations of Internet-based interventions for common mental disorders. *World Psychiatry* 2014 Feb;13(1):4-11 [FREE Full text] [doi: [10.1002/wps.20083](https://doi.org/10.1002/wps.20083)] [Medline: [24497236](https://pubmed.ncbi.nlm.nih.gov/24497236/)]
43. Titov N, Andrews G, Kemp A, Robinson E. Characteristics of adults with anxiety or depression treated at an internet clinic: comparison with a national survey and an outpatient clinic. *PLoS One* 2010 May 28;5(5):e10885 [FREE Full text] [doi: [10.1371/journal.pone.0010885](https://doi.org/10.1371/journal.pone.0010885)] [Medline: [20526371](https://pubmed.ncbi.nlm.nih.gov/20526371/)]

44. Hedman E, Ljótsson B, Andersson E, Andersson G, Lindefors N, Rück C, et al. Psychometric properties of Internet-administered measures of health anxiety: an investigation of the Health Anxiety Inventory, the Illness Attitude Scales, and the Whiteley Index. *J Anxiety Disord* 2015 Apr;31:32-37. [doi: [10.1016/j.janxdis.2015.01.008](https://doi.org/10.1016/j.janxdis.2015.01.008)] [Medline: [25723780](https://pubmed.ncbi.nlm.nih.gov/25723780/)]
45. Hoffmann D, Halsboe L, Eilenberg T, Jensen J, Frostholm L. A pilot study of processes of change in group-based acceptance and commitment therapy for health anxiety. *J Contextual Behav Sci* 2014 Jul;3(3):189-195 [FREE Full text] [doi: [10.1016/j.jcbs.2014.04.004](https://doi.org/10.1016/j.jcbs.2014.04.004)]
46. Pearcy CP, Anderson RA, Egan SJ, Rees CS. A systematic review and meta-analysis of self-help therapeutic interventions for obsessive-compulsive disorder: is therapeutic contact key to overall improvement? *J Behav Ther Exp Psychiatry* 2016 Jun;51:74-83. [doi: [10.1016/j.jbtep.2015.12.007](https://doi.org/10.1016/j.jbtep.2015.12.007)] [Medline: [26794856](https://pubmed.ncbi.nlm.nih.gov/26794856/)]
47. Berger T. The therapeutic alliance in internet interventions: a narrative review and suggestions for future research. *Psychother Res* 2017 Sep;27(5):511-524. [doi: [10.1080/10503307.2015.1119908](https://doi.org/10.1080/10503307.2015.1119908)] [Medline: [26732852](https://pubmed.ncbi.nlm.nih.gov/26732852/)]

Abbreviations

ACT: acceptance and commitment therapy
DSM-IV: Diagnostic and Statistical Manual of Mental Disorders, 4th edition
DSM-V: Diagnostic and Statistical Manual of Mental Disorders, 5th edition
HRQoL: health-related quality of life
iACT: internet-delivered acceptance and commitment therapy
iCBT: internet-delivered cognitive-behavioral therapy
MCS: mental component summary
PCS: physical component summary
RCT: randomized controlled trial
SCAN: Schedules for Clinical Assessment in Neuropsychiatry
SCL: symptom checklist
SF-12: short-form health survey
SRM: standardized response mean
WI-7: Whiteley-7 Index

Edited by J Torous; submitted 05.11.17; peer-reviewed by L McCracken, W Simm; comments to author 12.12.17; revised version received 05.02.18; accepted 14.02.18; published 06.04.18

Please cite as:

Hoffmann D, Rask CU, Hedman-Lagerlöf E, Ljótsson B, Frostholm L
Development and Feasibility Testing of Internet-Delivered Acceptance and Commitment Therapy for Severe Health Anxiety: Pilot Study
JMIR Ment Health 2018;5(2):e28
URL: <http://mental.jmir.org/2018/2/e28/>
doi: [10.2196/mental.9198](https://doi.org/10.2196/mental.9198)
PMID: [29625957](https://pubmed.ncbi.nlm.nih.gov/29625957/)

©Ditte Hoffmann, Charlotte Ulrikka Rask, Erik Hedman-Lagerlöf, Brjánn Ljótsson, Lisbeth Frostholm. Originally published in JMIR Mental Health (<http://mental.jmir.org>), 06.04.2018. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR Mental Health, is properly cited. The complete bibliographic information, a link to the original publication on <http://mental.jmir.org/>, as well as this copyright and license information must be included.